



THE HOSPITAL FOR
SICK CHILDREN

Department of
Pharmacy

baclofen 5 mg/mL Oral Suspension

Batch No: _____

Ingredients	Mfr	Lot #	Expiry Date	Quantity	Measured	Checked
baclofen 20 mg tablets	APO/Novartis			30		
Glycerin BP				4		
Simple Syrup NF				q.s. 120 mL		

Additional Information:

Need larger bottle to shake suspension. If settling on bottom of bottle cannot be shaken back into suspension then remove cap, stir with stirring rod, replace cap, shake bottle well, measure dose.

Equipment:

mortar and pestle glass stirring rod 5 mL oral syringe
graduated measure

Procedure:

Follow your Dept. procedures for risk assessment/training/PPE/equipment/facilities/NAPRA level


1. Crush tablets in the mortar to a fine powder with a pestle, or, soak tablets in glycerin and a small amount of vehicle for at least 120 minutes.
2. Add glycerin and a small amount of vehicle to crushed powder and levigate to a smooth paste with a pestle. soaked tablets, then levigate tablets into a smooth paste with a pestle. Continue to levigate as vehicle is added in small amounts until a liquid is formed.
3. Transfer liquid contents from mortar to graduate.
4. Use a small amount of vehicle to rinse mortar and add it to graduate.
5. Use vehicle to q.s. to the final volume. Stir well.
6. Transfer to amber bottle and label.

Quality Control:

Expected Product Appearance	Additional Notes
white suspension	

Storage: Refrigerate
Packaging: 250 mL amber glass bottle
BUD: 35 days

Sample Label:

 <p>THE HOSPITAL FOR SICK CHILDREN Department of Pharmacy</p>	baclofen 5 mg/mL Oral Suspension	
	Lot:	BUD:
	Refrigerate	Shake Well

Date Made/Prepared By/Checked By: _____

Reference:

1. Johnson CE. Stability of an extemporaneously compounded baclofen oral liquid. AJHP 1993;50:2353

Formulation Reviewed: October, 2020

Disclaimer: The information in this compounding and preparation worksheet was developed for in-house use only by The Hospital for Sick Children ("SickKids") and its staff. While SickKids endeavours to keep this information up to date and accurate, patient circumstances will vary and some information may become outdated as a result of recent medical developments following the time of printing. Any reliance you place on such information is therefore strictly at your own risk. Please note that the doses provided do not necessarily conform to the manufacturer's recommendations, and formulary drugs may not be licensed for use in children. The reader is cautioned to check the product information sheet provided by the manufacturer, and to verify indications, contraindications and recommended dosages. SickKids make no representations or warranties of any kind, express or implied, about the completeness, accuracy, reliability, or suitability of the information and related products contained in this worksheet for any purpose. SickKids accepts no responsibility for the reader's use, misuse or misinterpretation of the information, and does not promise success in achieving any desired result. In no event will SickKids be liable for any damage, liability or loss or for any indirect, incidental or consequential damages arising from the use of this information.